






















YOUR CHILD CAN STILL COME TO SCHOOL IF THEY HAVE:

A cough	A cold	A sore throat	Conjunctivitis	Headlice
			 Notify staff	 With treatment
Tonsillitis	Hand, Foot & Mouth	Headache	Tired	Tummy ache
	 Notify staff			
	Prescribed medication and non-prescribed medicine (e.g. paracetamol) can be administered to children, with parental consent.			

YOUR CHILD NEEDS TO STAY AT HOME IF THEY HAVE

<p style="text-align: center;">A temperature</p>  <p>To return to school when controlled by Calpol.</p>	<p style="text-align: center;">Sickness bug</p>  <p>To return to school after 48 hours of last episode.</p>	<p style="text-align: center;">Diarrhoea</p>  <p>To return to school after 48 hours of last episode.</p>	<p style="text-align: center;">Chicken pox</p>  <p>To return to school when the spots have crusted over.</p>	<p style="text-align: center;">Impetigo</p>  <p>To return to school when sores are crusted and healed or 48 hours after starting antibiotic treatment</p>
<p style="text-align: center;">Flu</p>  <p>Return to school when recovered/feeling better.</p>	<p style="text-align: center;">Slapped cheek</p>  <p>To return to school once the rash has developed.</p>	<p style="text-align: center;">Strep throat</p>  <p>To return to school 24 hours after starting antibiotics & feeling well enough</p>	<div style="display: flex; justify-content: space-between; align-items: center;">   </div> <p style="text-align: center;">REMEMBER!</p> <p style="text-align: center;">If your child is too ill to attend school, you MUST let school know as soon as possible.</p> <p style="text-align: center;">Please phone: 01709 760471</p>	